



2220 Pine View Way P.O. Box 4559 Petaluma, CA 94955-4559
 Telephone: (707) 765-6666 Fax: (707) 765-6674
 e-mail: info@scottlab.com website: www.scottlab.com



TANIN PLUS

Specific finishing tannin for white, red or rosé wines

Tanin Plus is a preparation of American oak tannins that has been lightly toasted to impart a slightly softer, more aromatic character. It also enhances the structure and organoleptic complexity of wine. Tanin Plus maintains the regular properties associated with the use of tannins such as helping with clarification, stability, oxidation, and aging.

RECOMMENDED DOSAGE

- 10-20g/hl for red wines (0.80-1.61lb/1000gal)
- 5-10g/hl for white wine s (0.40-0.80lb/1000gal)

Dissolve the tannin in 10 times its weight in warm water (35°C/95°F) and then add to the wine during a pumpover for good mixing.

Addition of Tanin Plus should be made a **minimum of three weeks** before bottling. After additions, proceed with normal racking.

PACKAGING

500g Bags

LABORATORY TRIAL PREPARATION

Prepare a 4.5% Tanin Plus solution by adding 4.5 grams of Tanin Plus to 80ml of deionized water in a 100ml volumetric flask. When dissolved, bring to volume of 100ml.

Laboratory Trial Recommendations:

Per 375 ml bottle

<u>Target Addition</u>	<u>lb/1000 gallon</u>	<u>mls of 4.5% Tanin Plus</u>
5 g/hl	0.41	0.41
10 g/hl	0.83	0.83
12 g/hl	1.00	1.00
14 g/hl	1.17	1.17
16 g/hl	1.34	1.34
18 g/hl	1.51	1.51
20 g/hl	1.66	1.66
30 g/hl	2.49	2.49

Allow wine samples to polymerize by waiting 12-24 hrs before tasting results.

The information herein is true and accurate to the best of our knowledge, however, this data sheet is not to be considered as a guarantee expressed or implied, or as a condition of sale of this product.